

# UCAC AND URAT: OPTICAL ASTROMETRIC CATALOG OBSERVING PROGRAMS

N. ZACHARIAS,<sup>1</sup> R. GAUME<sup>1</sup>

<sup>1</sup> U.S. Naval Observatory

3450 Mass.Ave. NW, Washington, DC 20392, USA

nz@usno.navy.mil

gaume@usno.navy.mil

**ABSTRACT.** The third USNO CCD Astrograph Catalog (UCAC3) all-sky catalog was released in August 2009. The final release, UCAC4, is in preparation and will include corrections and improvements to positions and proper motions, as well as publication of intermediate data. Properties of UCAC products are presented.

The USNO Robotic Astrometric Telescope (URAT) project uses the optics of the UCAC telescope with a new, wide-field camera to cover 28 sq.deg per exposure. A new all-sky survey to  $R \approx 18$  mag will begin in 2011 from Arizona and be continued from Chile. Neutral density spots on the single bandpass filter allow observations of bright stars. URAT will be able to directly link Hipparcos stars and extragalactic sources of the International Celestial Reference Frame (ICRF). Multiple sky overlaps per year will give mean positions, proper motions and parallaxes from these CCD observations. An accuracy level of 10 mas is expected for high signal-to-noise stellar observations.

## 1. INTRODUCTION

The goal of the astrometric observing programs described here is to provide many millions of accurate, all-sky, optical reference stars fainter than Hipparcos and Tycho-2 stars, at current epochs, including positions and proper motions. These star catalogs are aiming at a densification of the optical reference frame on the coordinate system of the defining radio frame (ICRF2, Fey et al. 2009). Due to errors in the proper motions of star catalogs the predicted positions at a future epoch degrade with time. Only new observations can improve the quality of the positions and proper motions.

name of catalog	ground space	proper motion	mag range	numb stars	pos.err (mas)	catalog release	remark
ICRF2	G	QSO	radio	3414	0.3	2009	VLBI
Hip.	S	yes	$\leq 12$	100 K	1.0	1997	ESA
Tycho-2	G/S	yes	$\leq 12$	2.5 M	10..100	2000	ESA,USNO
UCAC3	G	yes	8..16	100 M	20.. 70	2009	first CCD survey
2MASS	G	no	IR	500 M	90	2003	1 epoch
USNO-B	G	yes	12..21	1000 M	200	2003	Schmidt plates
PanSTARRS	G	yes	17..23	2000 M	$\leq 30$	$\geq 2011$	Hawaii
SkyMapper	G	yes	16..22	1000 M	?	$\geq 2011$	Australia
SST	G	yes	17..23	2000 M	?	$\geq 2011$	DARPA
URAT1	G	yes	8..18	500 M	10..30	2012	USNO,NOFS,CTIO
LSST	G	yes	18..24	3000 M	?	$\geq 2015$	NOAO
nano-JASMINE	S	yes	near IR	2 M	3.0	$\geq 2013$	Japan
JMAPS	S	yes	0..14	30 M	1.0	2018	USNO
Gaia	S	yes	6..20	1000 M	0.025	$\geq 2015$	ESA
SIM	S	yes	0..20	20,000	0.004	on hold	NASA,JPL

Table 1: Comparison of current and future (nearly) all-sky survey observing projects significant for astrometry.

Table 1 provides an overview about current and future, astrometric, global catalog observing programs

from the ground and space. The UCAC program is essentially complete, pending a final data release, while the URAT program is a ground-based follow-up to UCAC to begin in 2011 with improved accuracy and reaching even fainter magnitudes with first results expected to be released in 2012. Contrary to other ground-based sky survey projects, URAT uses a small-aperture, dedicated astrometric telescope for highest positional accuracy of stars in the mid-magnitude range, bridging the gap between the current optical reference frame and deeper surveys.

## 2. UCAC

The USNO CCD Astrograph Catalog project observations were completed in 2004. Operating in a bandpass between V and R, over 278,000 exposures of 1 sq.deg each were obtained in a 2-fold all-sky overlap pattern with each field having a long (125 sec) and a short (25 sec) exposure. After the UCAC1 early data release (Zacharias et al. 2000), the widely used UCAC2 catalog (Zacharias et al. 2004) was released, covering declinations  $-90^\circ$  to about  $+50^\circ$  for over 48 million stars.




Figure 1: Precision of re-reduced NPM first epoch data. The standard error for the  $RA \cdot \cos(Dec)$  coordinate is shown as a function of NPM blue magnitude (Girard 2010). Results for the Dec coordinate are similar.

UCAC2 is a relatively “clean” reference star catalog, having cut out all identified “problem” cases (like blended images), while the UCAC3 released in 2009 is the first all-sky catalog in this series (Zacharias et al. 2010), aiming at a high degree of completeness and thus also includes a relatively large number of problem cases and unreliable low S/N sources. Details about the astrometric reductions of UCAC3 can be found elsewhere (Finch et al. 2010).

UCAC3 was put together within a given deadline and lacks high-quality proper motions for faint stars north of declination  $-30^\circ$ . Uncorrected SuperCosmos data of Schmidt plate scans were widely used. This resulted in systematic errors in UCAC3 proper motions as function of the Schmidt plate pattern of up to about 10 mas/yr in some areas (Zacharias et al. 2010, Röser et al. 2010).

Most stars brighter than about  $V=12.5$  and many others brighter than  $V=14$  have high-quality proper motions in UCAC3 due to the use of early epoch astrograph plates. Over 5000 plates from the AGK2 (epoch close to 1931), as well as from the Hamburg Zone astrograph and USNO Black Birch twin-astrograph programs, have been scanned on the StarScan machine (Zacharias et al. 2008) and reduced for UCAC. All other faint stars to the limit of UCAC and south of  $\delta = -30^\circ$  also have high-quality proper

motions thanks to the Southern Proper Motion (SPM) program and re-reductions in a joint USNO/Yale effort, which provided SPM 1st epoch data in time for the UCAC3.

The proper motion problems of UCAC3 north of  $\delta = -30^\circ$  and software bugs causing for example some stars (about 1% of all stars in the catalog) to be listed twice and others not at all, will be resolved in the upcoming, final, UCAC4 release. In a collaboration between USNO and Yale University, the Northern Proper Motion (NPM) Lick astrograph plates (which were also scanned on the PMM machine at NOFS, like the SPM plates) were re-processed to provide a new, highly accurate star catalog at a mean epoch of about 1950 covering all sky north of the SPM data. Fig. 1 shows the random errors of the NPM 1st epoch data and combined with the  $\approx 50$  year epoch difference to UCAC CCD observing, proper motion errors of 3 to 5 mas/yr are expected with remaining systematic errors estimated to be on the 2 mas/yr level. In addition to a compiled catalog of about 100 million stars, individual CCD observed positions will become available with the UCAC4 release in 2011.

### 3. OPTICAL REFERENCE FRAME TODAY

About 20 years after the central epoch of observations the positional accuracy of individual Hipparcos stars has degraded to typically 20 mas per coordinate ( $1 \sigma$ ). The coordinate axes of the Hipparcos reference frame are still aligned to the ICRF with errors not larger than 2.7 mas and system rotations not larger than 0.55 mas/yr; limits set by a recent investigation of 46 radio stars (Boboltz et al. 2007).




Figure 2: 2-dim (RA,Dec) position discrepancies of selected Hipparcos stars. A red, open circle represents the new Hipparcos reduction position, while a green, filled circle shows the original Hipparcos catalog position relative to that and the blue diamond indicates the location of the UCAC CCD observation as obtained near the 2000 epoch. The respective Hipparcos catalogs position, proper motion and parallax data was used to derive a position at the UCAC epoch for this comparison. The size of the box for each star is 1 arcsec = 1000 mas.

However, some Hipparcos stars can have much larger positional errors at current epochs. During the UCAC3 construction about 1500 stars (random sample) were excluded as reference stars and the observed CCD-based positions (at about epoch 2000) compared with the original (ESA 1997) and re-reduced (van Leeuwen 2007) Hipparcos catalog positions at that epoch. For about 1 to 2 % of the stars large discrepancies were found (Zacharias et al. 2009) up to several 100 mas (Figure 2). Sometimes the CCD data agree with the original but not the new reductions and vice versa, and sometimes all 3 data

points are not consistent with estimated errors by large margins.

Limits of the Tycho-2 reference frame begin to become visible as well. Systematic radio-optical extragalactic reference frame source position differences are seen in the 30 mas range for some areas, which may be due to remaining Tycho-2 systematic errors (Zacharias & Zacharias 2009). Small magnitude equations of Tycho-2 positions are seen in UCAC3 data (Zacharias et al. 2010) and in work with the SPM (Girard, private communication) as function of declination zones.

Going to fainter stars, systematic errors of star positions at current epochs are at least 20 mas just from the expected error propagation of proper motion errors in UCAC type catalogs. For Schmidt plate data, these local systematic errors reach 200 to 400 mas, as seen in many external comparisons.

#### 4. URAT

The USNO Robotic Astrometric Telescope (URAT) project uses the same astrograph “red lens” as was used for the UCAC program. A completely new tube assembly has been built in the USNO Instrument Shop and electronic upgrades of its B&C mount are in progress. For the first time, on September 9, 2010, the new mount system was operated under computer control to point at stars and track.




Figure 3: The new tube assembly of the USNO astrograph in 2009 after completion of the mechanical work. The camera dewar has a mass of  $\approx 140$  kg and the change in mass due to evaporating  $LN_2$  requires a movable counterweight on the opposite side to balance this relatively small telescope on its mount.

telescope aperture	20	cm
bandpass	670–750	nm
pixel scale	0.905	arcsec/pixel
pixel size	9.0	$\mu\text{m}$
field of single detector	2.65 x 2.65	degree
number of imaging detectors	4	STA1600
number of guide/focus det.	3	2k by 5k each
sky area per exposure	28	sq. degree
survey expos.time	4	min
magnitude range	8 – 18	mag
survey begin at NOFS	2011	

Table 2: Properties of the URAT program.

A summary of the URAT project properties is given in Table 2. The detector design and electronics allow for clocked anti-blooming operations which extend the usable dynamic range for astrometry by about 3 magnitudes at the bright end (beyond saturation). The dewar has a 300 mm clear aperture window which also serves as the filter. It contains also 2 neutral density spots (5 and 7.5 mag attenuation, respectively) to allow stars as bright as about  $R = 1$  be observed relative to surrounding stars of magnitude 8 or fainter in a large area of sky. A prototype camera with a single, front-side illuminated 10k detector was successfully operated in 2007/2008. A “4-shooter” camera for URAT was funded in 2008. The 4 thinned STA16000 detectors of 10,560 by 10,560 pixels each (Figure 3) image 28 square degrees of sky in a single exposure. On 3 sides of the 2 by 2 array of main detectors, there are additional 2k by 5k CCDs with 8  $\mu\text{m}$  pixel size used for guiding and focus control.



Figure 4: Flat field image of a successfully thinned 10k detector for the URAT camera. Pixel data of the 16 outputs are shown with overscans. The cosmetic quality is excellent with only 2 column defects on the entire 111 million pixel area.

After test and commissioning the entire instrument will be moved to the USNO Flagstaff, Arizona station (NOFS) in 2011 to begin a new sky survey. The shipping container also serves as control room.

After 2 to 3 years of operation, the instrument will likely be moved to the Cerro Tololo Interamerican Observatory (CTIO) to observe the southern hemisphere. Due to the large field of view, several complete sky overlaps per year will be obtained of the accessible hemisphere to derive mean positions, proper motions and parallaxes on the 10 to 30 mas level, depending on the brightness of stars.

The URAT survey will be able to directly link many ICRF optical counterparts to the Hipparcos reference frame. A successful Gaia mission of course will supersede most of the URAT data; however, a significant improvement of the optical reference frame and data for galactic dynamics studies are expected from URAT in several releases beginning in 2012. URAT will also provide excellent reference stars for the other ongoing or planned very deep surveys until then.

For the latest update on UCAC and URAT, see <http://www.usno.navy.mil/usno/astrometry/optical-IR-prod>.

## 5. REFERENCES

- Boboltz, D. A., Fey, A. L., Puatua, W., Zacharias, N., Claussen, M. J., Johnston, K. J., Gaume, R. A., 2007, “VLA+PT Astrometry of 46 Radio Stars” *AJ* 133, pp. 906
- ESA 1997, “The Hipparcos Catalogue”, ESA SP-1200
- Fey, A., Gordon, D., Jacobs, C.S. (Eds.), 2009, “The second realization of the international celestial reference frame by very long baseline interferometry”, IERS Technical Note No. 35, Frankfurt
- Finch, C., Zacharias, N., Wycoff, G., 2010, “UCAC3 astrometric reductions”, *AJ* 139, 2200–2207
- Girard, T., 2010, private communication.
- Röser, S., Demleitner, M., Schilbach, E., 2010, “The PPMXL catalog of positions and proper motions on the ICRF”, *AJ* 139, pp. 2440
- van Leeuwen, F. 2007, “Hipparcos, the new reduction of the raw data”, *A&A* 474, pp. 653
- Zacharias, M.I., Zacharias, N., 2009, “Significant radio-optical reference frame offsets from CTIO data”, poster paper IAU General Assembly, Rio de Janeiro
- Zacharias, N., Urban, S.E., Zacharias, M.I., Hall, D.M., Wycoff, G.L., Rafferty, T.J., Germain, M.E., Holdenried, E.R., Pohlman, J.W., Gauss, F.S., Monet, D.G., Winter, L., 2000, “The first US Naval Observatory CCD Astrograph Catalog”, *AJ* 120, pp. 2131
- Zacharias, N., Urban, S.E., Zacharias, M.I., Wycoff, G.L., Hall, D.M., Monet, D.G., Rafferty, T.J., 2004, “The Second US Naval Observatory CCD Astrograph Catalog (UCAC2)”, *AJ* 127, pp. 3043–3059
- Zacharias, N., Winter, L., Holdenried, E.R., De Cuyper, J.-P., Rafferty, T.J., Wycoff, G.L., 2008, “The StarScan plate measuring machine: overview and calibrations”, *PASP* 120, pp. 644
- Zacharias, N., Finch, C., Wycoff, G., Hartkopf, W., 2009, “Improving Hipparcos Proper Motions with UCAC”, DDA meeting, Virginia Beach, *BAAS* 41 No.2, p. 910, abstract 16.03
- Zacharias, N., Finch, C., Girard, T., Hambly, N., Wycoff, G., Zacharias, M.I., et al. 2010, “The Third US Naval Observatory CCD Astrograph Catalog (UCAC3)”, *AJ* 139, pp. 2184–2199